Cleaning & Disinfection Protocol for Non-Enveloped Viruses

This document has been developed in accordance with current applicable infection control and regulatory guidelines. It is intended for use as a guideline only. At no time should this document replace existing documents established by the facility unless written permission has been obtained from the responsible facility manager.

PREFACE

Non-enveloped viruses lack a lipid-bilayer membrane. To reproduce they must breach the membrane of a target host cell to gain access to its cytoplasm. The mode of transmission for non-enveloped viruses is characterized by the specific virus; however, the most common routes are via indirect or direct contact of infectious virus particles, contact with or inhalation of respiratory droplets. Some non-enveloped viruses may also be spread via airborne transmission. Unlike, enveloped viruses, non-enveloped viruses are not easily inactivated by many routine surface cleaning and disinfectant products.

The following table provides examples of Non-Enveloped Viruses and Mode of Transmission of concern for Healthcare settings.

Virus	Mode of Transmission	Infective Material
Adenovirus	Large droplets; direct and	Respiratory Secretions
	indirect contact	
Astrovirus	Direct and indirect contact	Feces
	(fecal/oral)	
Norovirus	Direct and indirect contact	Feces
	(fecal/oral)	
Chickenpox	Direct and indirect contact;	Lesion drainage, Respiratory
	Airborne	Secretions
Coxsackievirus	Direct and indirect contact	Feces, Respiratory Secretions
Echovirus	Direct and indirect contact	Feces, Respiratory Secretions
Enterovirus	Direct and indirect contact	Feces, Respiratory Secretions
Hepatitis A	Direct and indirect contact	Feces
	(fecal/oral)	
Hepatitis E	Direct and indirect contact	Feces
	(fecal/oral)	
Parvovirus B-19	Direct contact; Large Droplets	Respiratory Secretions
Poliovirus	Direct and indirect contact	Feces, Respiratory Secretions
Rhinovirus	Direct and indirect contact;	Respiratory Secretions
	possibly large droplets	
Rotavirus	Direct and indirect contact	Feces



Cleaning & Disinfection Protocol for Non-Enveloped Viruses

PREPARATION

Transmission of non-enveloped viruses can be attributed to direct and indirect contact, respiratory droplets and airborne transmission. Appropriate personal protection should be taken for those responsible for the decontamination of a room or area. Appropriate bio-security practices should be applied, including limiting the amount of aerosols generated and disturbance to dust / soil in the area to be cleaned and disinfected.

PROTECTIVE BARRIERS

Appropriate personal protection should be taken for those responsible for the decontamination of a room or area.

- 1. Disposable gloves. Gloves should be changed as required; i.e., when torn, when hands become wet inside the glove or when moving between patient areas.
- 2. Protective Eye wear (goggles, face shield or mask with eye protection), as needed
- 3. Masks (surgical or procedural masks sufficient), as needed
- 4. Gowns, as needed

PRODUCTS

All disinfectant or disinfect-cleaner products to be used for cleaning and disinfection of environmental surfaces and patient care equipment must be approved by either the Environmental Protection Agency (EPA) or Health Canada and carry an EPA or Drug Identification Number (DIN). Products claiming to be a disinfectant but do not carry an EPA number/DIN have not been approved for sale in the US or Canada and should not be used. Non-enveloped viruses are in the same category as Mycobacteria with respect to difficulty of inactivation, therefore, it is important to review the product label for efficacy against non-enveloped viruses. Choose products that either have proven efficacy against the virus of interest or that carry the Broad-Spectrum Virucide Claim (in Canada only) which in accordance to Health Canada proves efficacy against both enveloped and non-enveloped viruses. In the US, look for products with proven efficacy against a number of enveloped and non-enveloped viruses.

Disinfectant Chemistries Approved for Disinfection of Environmental Surfaces & Patient Care Equipment include:

- 1. Accelerated Hydrogen Peroxide® (AHP®)
- 2. Sodium Hypochlorite
- 3. Quaternary Ammonium Compounds
- 4. Alcohol
- 5. Phenolics



2770 Coventry Road Oakville, Ontario L6H 6S2 Tel: 1-800-387-7578 Fax: (905)813-0220 www.infectionpreventionresource.com

Cleaning & Disinfection Protocol for Non-Enveloped Viruses

The concentration and contact time for each product will differ. For that reason it is important to read the product label prior to commencing any cleaning and disinfection process.

RECOMMENDED PROCEDURES FOR CLEANING AND DISINFECTION

Summary of Procedure: Apply the solution to either the surface or device surface or to cloth. Clean all horizontal surfaces in the room ensuring that the cloth is changed when soiled. Appropriately dispose of cloth. Disinfect all horizontal surface of the room by applying the disinfectant and allowing for contact time as per the product label. Allow surfaces to air dry or wipe dry with a clean, dry cloth if surfaces are still wet after the contact time as been achieved. Periodic rinsing of soft surfaces such as vinyl or naugahyde is suggested as well as equipment regularly handled by hand. Bathrooms within a room should be cleaned last.

- 1. Gather all equipment, cleaning solutions and materials required to clean the environmental surfaces.
- 2. Wash hands and put on gloves prior to cleaning the surface. Personal protective equipment should be changed if torn or soiled.
- 3. Visible or gross soil present and/or blood or body fluid spills must be removed prior to cleaning. [See Protocol for Cleaning & Disinfecting a Blood or Body Fluid spill.]
- 4. As appropriate clean all high touch clinical contact and housekeeping surfaces. To ensure that cross contamination does not occur use clean cloths for each surface being cleaned. If using an open bucket system, ensure that solutions do not become contaminated (NO DOUBLE DIPPING). If using a disposable wipe system, ensure that a new wipe is being used for each surface being cleaned.
- 5. To disinfect environmental surfaces, apply the disinfectant and allow surfaces to remain wet for the appropriate contact time as specified on the product label.
- 6. If using a 1-Step Cleaning-Disinfecting Solution a separate cleaning step is not necessary unless the surfaces are visibly soiled. To ensure disinfection occurs, the cleaner-disinfectant solution may need to be applied multiple times in order to achieve the contact time as specified on the product label.
- 7. Soiled rags should be placed in a bag for laundering. Disposable cloths should be disposed as regular waste in garbage bags or as directed on product label.
- 8. Remove and discard gloves
- 9. Wash hands

RECOMMENDED PROCEDURES FOR CLEANING AND DISINFECTION OF BLOOD AND BODY FLUIDS

Appropriate personal protective equipment should be worn for cleaning up a body fluid spill. Gloves should be worn during the cleaning and disinfecting procedures. If the possibility of splashing exists, the worker should wear a face shield and gown. For large spills, overalls, gowns or aprons as well as boots or



Cleaning & Disinfection Protocol for Non-Enveloped Viruses

protective shoe covers should be worn. Personal protective equipment should be changed if torn or soiled, and always removed before leaving the location of the spill, and then wash hands.

- 1. Wash hands and put on gloves.
- 2. If the possibility of splashing exists, the worker should wear a face shield and gown. For large spills, overalls, gowns or aprons as well as boots or protective shoe covers should be worn. Personal protective equipment should be changed if torn or soiled and always removed before leaving the location of the spill.
- 3. Apply the disinfectant solution to spill wait until the contact time has been achieved.
- 4. Blot up the blood with disposable towels. Dispose of towels in plastic-lined waste receptacle.
- 5. Spray or wipe surface with the disinfectant solution ensure the appropriate contact time is achieved. Wipe dry with disposable paper towel. Discard paper towel as above.
- 6. Remove gloves and dispose in plastic-lined waste receptacle.
- 7. Wash hands.

DISPOSAL OF INFECTIOUS MATERIAL

All cleaning cloths gloves and handled tools used for the decontamination of suspected infectious diseases must be placed in a clearly marked plastic lined waste receptacle. Decontaminate all wastes before disposal; i.e., steam sterilization, chemical disinfection and or incineration.

REFERENCES

APIC, Ready Reference To Microbes, 2002

Best Practices for Cleaning, Disinfection and Sterilization in All Health Care Settings, Provincial Infectious Diseases Advisory Committee (PIDAC), May 2013

Best Practices for Environmental Cleaning for Prevention and Control of Infections in All Health Care Settings, PIDAC, May 2012

Guidelines for Environmental Infection Control in Healthcare Facilities, CDC. MMWR June 2003, Vol 52, No RR-10

Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings, HICPAC, 2007

Rutala WA & Weber DJ. The benefits of surface disinfection. AJIC 2004;32(4) 226-229



Cleaning & Disinfection Protocol for Non-Enveloped Viruses

Mayhall CG. Hospital Epidemiology and Infection Control, 3rd Ed. Philadelphia. Lippincott Williams & Wilkins, 2004

